

LIBRARY
College of Dentistry
UNIVERSITY OF SOUTHERN CALIFORNIA

The
**DENTAL
DIGEST**



SEE PAGE 166

THE FIRST HUNDRED YEARS

AUGUST, 1936
Vol. 42 No. 8



"and Doctor ...
**YOU'VE MADE ME CHANGE MY
 MIND ABOUT DENTAL WORK"**

● You appreciate the possibilities offered you through the use of local anesthesia in obtaining the confidence of your patients. And, the selection of any anesthetic solution (Cook) can play an important part in gaining their acceptance of local anesthesia.

Novocain-epinephrin solutions A and AB possess the essentials which promote patient-comfort during and after your operations. Novocain with Cobefrin offers you more. The use of Cobefrin as the vasoconstrictor in this solution offers a higher degree of patient-tolerance in normal and abnormal patients alike.

Note: It is the use of a normal saline vehicle that anesthetic solutions (Cook) differ from those available in R. B. Waite cartridges and ampules.

Cobefrin, Trademark Reg. U. S. Pat. Off. and Canada.
 Winthrop Chemical Co., Inc., Brand of Nordofrin.

Carpule
REG. U.S. PAT. OFF. & CANADA



COOK

COOK LABORATORIES, INC.

170 Varick Street, New York, N. Y. . . . Laboratories: Rensselaer & Springville, N. Y.

1891

1936



Distinctive
ANESTHETIC SOLUTIONS..

● Anesthetic solutions (R. B. Waite) for 45 years have been a byword for anesthetic efficiency and safety in dental offices throughout the world. During this period, many improvements have been made in these solutions and in the manner in which they have been supplied to the profession.

Today, Novocain-epinephrin solutions 2 and 4 and Novocain with Cobefrin (R. B. Waite) offer the utmost in uniformity, efficiency and safety. At the same time they offer the same intrinsic qualities that have always enjoyed the confidence of leading practitioners everywhere . . . the same compound base which is their distinguishing feature.

If you are interested in learning more about the peculiar advantages of this base, ask your salesman or write this office.


Novocain, Trademark Reg. U. S. Pat. Off. and Canada.
 Winthrop Chemical Co., Inc., Brand of Procaine HCl.



R.B. Waite
REG. U.S. PAT. OFF. & CANADA

R.B. WAITE

The ANTIDOLOR MFG. CO., INC.



Who *wouldn't* feel Happy!...

HE BEGAN wearing his new denture only yesterday, yet he's already on accustomed terms with his ever enjoyable cigar. That seems to him a veritable feat; his dentist, however, had merely prescribed DR. WERNET'S Powder.

It forms an elastic, adhesive, and protective cushion between the denture and the tissues. It holds the denture, *mechanically*, more firmly in

place, and soothes and protects tissues that are tender. By promoting comfort and greater assurance, it makes it possible for the denture to be worn more regularly, thus hastening its mastery by the patient.


Send for YOUR supply—FREE!

Simply mail the lower portion of this page with your card or letterhead to WERNET DENTAL MFG. CO., 882 Third Ave., Brooklyn, N. Y.

DR. WERNET'S POWDER

Speeds the Mastery of a New Denture!

When Treating VINCENT'S ANGINA



Restrain Systemic Disease with SAL HEPATICA

Sal Hepatica can obviate many systemic complications which occur so often with Vincent's Angina and other foci through (1) rapid and effective removal of intestinal waste (2) neutralization of acidity and maintenance of a normal alkaline level—an excellent offensive against disease.

In components, their ratio and action, Sal Hepatica approximates the famous natural saline waters. It makes an agreeable, effervescent drink.

***SAL HEPATICA CLEANS
THE INTESTINAL TRACT
AND COMBATS ACIDITY***

Sample on Request

BRISTOL-MYERS CO.

19-S West 50th Street
New York, N. Y.

(Please use coupon on page 286)

The DENTAL DIGEST

VOLUME 42

August, 1936

NUMBER 8

Extra-Oral Roentgenology	- - - - -	260
<i>George W. Matthews, B.S., D.D.S., M.S.</i>		
Administration of Ethyl Chloride Anesthesia	- - - - -	264
<i>Julius A. Greenhouse, D.D.S.</i>		
"The First Hundred Years"	- - - - -	266
<i>Joseph J. Tolan, D.D.S., M.S.</i>		
Restoring the Upper First Bicuspid	- - - - -	270
<i>Jerome M. Schweitzer, D.D.S.</i>		
Cast Individual Impression Trays	- - - - -	272
<i>Leonard S. Fletcher, D.D.S.</i>		
Stone Models for Indirect Castings	- - - - -	274
<i>Robert S. Hines, D.D.S.</i>		
The Editor's Page	- - - - -	275
A Rebasing Technique	- - - - -	276
<i>P. N. Ogle, D.D.S.</i>		
About Our Contributors	- - - - -	278
Suggestions to Contributors	- - - - -	288

EDWARD J. RYAN, B. S., D. D. S., Editor

ETHEL H. DAVIS, A. B., Assistant Editor

AN ORAL HYGIENE PUBLICATION. Published monthly on the fifteenth by Dental Digest, Inc. Copyright, 1936, by Dental Digest, Inc. Entered as second class matter at the Postoffice at Ashland, Ohio, under the Act of Congress, March 3, 1879. PUBLICATION OFFICES: 1005 Liberty Avenue, Pittsburgh, Pennsylvania. Merwin B. Massol, Publisher; Associates: J. J. Downes, W. Earle Craig, D. D. S.; R. C. Ketterer, Publication Manager. Subscriptions should be sent to the Publication Offices, 1005 Liberty Avenue, Pittsburgh, Pennsylvania. Manuscripts and correspondence regarding editorial matters should be addressed to the editor at 708 Church Street, Evanston, Illinois. Subscription, including postage: \$2 per year in the United States, Alaska, Cuba, Guam, Hawaiian Islands, Mexico, Philippines, Puerto Rico. To Great Britain and Continent, \$2.75; Canada, \$2.00; Australia, \$2.75. All other countries, \$2.75. Single copies, 25c. DISTRICT ADVERTISING OFFICES: *Chicago*: Peoples Gas Building; W. B. Conant, Western Manager. *New York*: 18 East 48th Street; Stuart M. Stanley, Eastern Manager. *St. Louis*: Syndicate Trust Building; A. D. McKinney, Southern Manager. *San Francisco*: 155 Montgomery Street. *Los Angeles*: 318 West 9th Street; Don Harway, Pacific Coast Manager.

EXTRA-ORAL ROENTGENOLOGY

GEORGE W. MATTHEWS, B.S., D.D.S., M.S.

Birmingham, Alabama


DENTAL roentgenologists who are not making use of extra-oral views of the jaws are not availing themselves of one of the best diagnostic aids at our disposal. Information can be obtained regarding structures which cannot

possibly be seen in intra-oral views. Extra-oral roentgenograms are also valuable when used in conjunction with intra-oral ones to show better relationship of structures. The operator can orient himself much better when the entire jaw is shown on one

plate, rather than when several small roentgenograms are used.

INDICATIONS

Extra-oral roentgenograms are invaluable in oral surgery and in diagnosis of an obscure condition or re-



TECHNIQUE FOR SYMPHYSIS OF THE MANDIBLE (Figs. 1, 2, 3, 4)

Fig. 1—Head and Cassette Position—Patient seated sidewise in chair, cassette resting on head-rest at angle of 45° with the floor, and steadied with patient's hand on lower corner. Patient's head is thrown backward as far as is comfortably possible and the chin and nose pressed firmly against the cassette, so that the cassette forms an angle of 45° with the sagittal plane of the head.

Figs. 2 and 3—Position and Direction of Tube—Tube is positioned under, behind, and slightly median to the angle of the jaw and directed so that the central ray will hit the cassette at a mesio-distal angle of 90° and a vertico-horizontal angle of about 70° . The central ray will pass between the cervical vertebrae and the ramus opposite to the side being roentgenographed, and should be directed at the lower cuspid region.

Fig. 4—Roentgenographic Anatomy—A clear view is obtained of the mandible from the second molar region to the median line. The roots of all teeth can be seen and also the cancellous bone and cortical plate of inferior border. The cervical vertebrae are superimposed over the inferior border of the mandible below the molars. The radiopaque area at the lower border of the mandible directly below the bicuspid is the hyoid bone.

This view is important in early diagnosis of osteomyelitis which frequently begins in the region of the mental foramen, necrosis originating from the anterior and bicuspid teeth, and fractures in the anterior part of the mandible.



flex pain in the jaws. Among the more common conditions disclosed by a careful and systematic extra-oral examination are: cysts, fractures, osteomyelitis, necrosis, odontomas, impacted, unerupted, and supernumerary teeth, and foreign bodies.

Although it must be emphasized that extra-oral roentgenograms are an adjunct to intra-oral ones there are unusual cases in which they are actually substituted in place of intra-oral views. These are:

1. Cases of trismus of the jaws in which a film cannot be placed and held inside the mouth.

2. Swelling and edema of the floor

of the mouth, preventing placing of a film packet.

3. Patients who gag violently when a film touches the soft palate despite all the standard preventives.

4. Lower third molar cases in which the film packet cannot be placed far enough posteriorly to include a horizontal or mesio-angular impaction owing to the posterior edge of the film striking the pterygoid muscles and the pharyngopalatine arch. There are many cases of this type.

5. Very young children.

It must be borne in mind that the field of extra-oral roentgenology covers a great deal more than lateral

views of the jaws. There are lateral views of the skull, anteroposterior views of the skull, special views of the temporomandibular articulation, and the various views for diagnosing conditions of the paranasal sinuses. In this article the lateral views of the jaw will be discussed. There are four variations of this view: (1) view of the symphysis of the mandible; (2) view of the body of the mandible; (3) view of the angle of the mandible, and (4) view of the ramus of the mandible.

INTERPRETATION

Interpretation of lateral jaw roent-


TECHNIQUE FOR THE BODY OF THE MANDIBLE (Figs. 5, 6, 7, 8)

Fig. 5—Head and Cassette Position—Patient seated sidewise in chair, cassette resting on head-rest at an angle of 45° with floor, and steadied with patient's hand on lower corner. Patient's head is thrown backward as far as is comfortably possible and the cheek pressed firmly against the cassette at such an angle that the tip of the nose is about an inch from the cassette.

Figs. 6 and 7—Position and Direction of Tube—Tube is under the angle and directed so that the central ray will hit the cassette at a mesio-distal angle of 90° and a vertico-horizontal angle of 70°. The central ray will pass between the cervical vertebrae and the ramus of the side opposite to that being roentgenographed, and should be directed at the first molar region.

Fig. 8—Roentgenographic Anatomy—A clear view is obtained of the body of the mandible from the third molar to the cuspid region. A clear profile view of the lower bicuspid and molar teeth is also shown but a good profile view of the uppers is not obtained owing to the fact that the rays do not pass directly through the contact points of these teeth; some overlap and distortion occurs. The mandibular canal can be traced just inferior to the roots of the molar teeth and the mental foramen can be seen directly between the apexes of the bicuspids. The radiopaque area at the lower border of the mandible directly under the first molar is the hyoid bone, and the cervical vertebrae are superimposed on the ramus.

Incidental Features—The upper and lower third molars are impacted and there is apparently some pericoronar infection of the lower third molar. There is apparently decay under the restoration in the lower first molar.



genograms requires a thorough knowledge of the normal anatomy of the head and particularly of the neck. The superimposition of structures must be taken into consideration and due allowance made for it. Radiopaque structures, through which the rays pass between the tube and the jaws, cast radiopaque areas over the image of the jaws on the plate; likewise, radiolucent structures, through which the rays pass between the tube and the jaws, cast radiolucent shadows on the plate. Both of these normal anatomic conditions have been misinterpreted as pathologic conditions.

The most commonly superimposed

structures are: (1) the maxilla and mandible of the side nearest the tube superimposed on the anterior part of the side being roentgenographed; (2) the cervical vertebrae; (3) the hyoid bone; (4) the cricoid cartilage; (5) the pharynx; (6) the tongue; (7) the palate and uvula, and (8) the zygomatic arch.

In addition to recognizing these normal anatomic conditions the oral diagnostician must, if possible, recognize and interpret the various pathologic lesions in their incipience. The classic roentgenographic appearance of osteomyelitis in its early stages should be known and instantly recognized; fine line fractures in obscure

regions, such as the head of the condyle, should be detected; necrosis and sequestrums should be recognized, and foreign bodies should be located by views from different angles.

GENERAL TECHNIQUE

1. All four variations of lateral jaw roentgenograms are best made in a dental chair, with the patient seated sidewise and the cassette resting on the head-rest at about a forty-five degree angle with the floor.

2. With an intensifying screen and super-speed film, an exposure of one and one-half seconds is usually all that is needed with the average den-



TECHNIQUE FOR ANGLE OF THE MANDIBLE (Figs. 9, 10, 11, 12)

Fig. 9—Head and Cassette Position—Patient seated sidewise in chair, cassette resting on head-rest at an angle of 45° with the floor. Patient's head thrown backward as far as is comfortably possible and the cheek pressed firmly against the cassette so that the ramus of the mandible lies flat against it. The tip of the nose will be about 1½ inches from the cassette. The patient steadies the cassette by holding the lower corner.

Figs. 10 and 11—Position and Direction of Tube—The tube is positioned under the angle of the jaw and directed so that the central ray will hit the cassette at a mesio-distal angle of 90° and a vertico-horizontal angle of 70°. The central ray will pass through the third molar region of the side being roentgenographed.

Fig. 12—Roentgenographic Anatomy—A clear view is obtained of the inferior border of the mandible from the mental foramen region back to and including the angle. The body of the opposite side is superimposed over all the teeth except the molars in this view. The pharynx is superimposed over the ramus of the mandible which accounts for the radiolucent appearance. The radiopaque area seen on the anterior half of the ramus is caused by the division of the pharynx into its oral and nasal parts. The radiopaque area in the pharynx just inferior to the angle is part of the hyoid bone. The mandibular canal can be seen running through the body of the mandible right at the third molar roots and well down below the roots of the other molars. The radiolucent area in the posterior half of the ramus is caused by the mandibular fossa and foramen.

Incidental Features—There is a fracture of the mandible just distal to the third molar which has virtually united. The mesial root of the lower third molar is fractured. This tooth was left in place until after the fracture was united to prevent the posterior fragment from rising. There is considerable periodontal involvement of both upper and lower molars. Calculus can be seen at the necks of all these teeth. The radiopaque area just distal to the lower third molar which has the appearance of a root is merely contrast between the radiolucency of the anterior wall of the pharynx and the line of fracture.



tal machine. It has not been found necessary to secure the patient's head to the cassette with a bandage for stability. This method is recommended by such authorities as Ennis and Simpson, and no doubt is essential in anteroposterior views when eight seconds and longer are required; but if the cassette is placed firmly on the head-rest with the patient's face firmly against it, and the cassette steadied with the patient's hand on the lower corner—then there is no movement.

3. The patient is instructed to hold his breath just prior to exposure, and also not to swallow. Swallowing during exposure causes an abnormal radiolucency of the pharynx.

PROCESSING

As stated the best results are obtained by using a cassette with intensifying screens. These screens must be kept scrupulously clean and no dirt, dust, or drops of water allowed on them. In opening the cassette and removing the film, care must be exercised not to slide the film across the screen, as friction produces static electricity which sometimes shows on the roentgenogram in the form of something that looks like a bolt of lightning.

The best results are obtained by developing at 65° F. for five minutes. If this produces a roentgenogram that

is too dark the exposure time should be reduced.

TARGET-FILM DISTANCE

It will be noted in the accompanying illustrations that the point of the cone is either touching or near the patient's face. This is only done in order to include the cone in the illustration. In actual practice much better results are obtained by backing up the tube to at least 18 inches or 2 feet. This gives less distortion and much clearer definition. Of course, the time will have to be increased accordingly, but this can be determined by each operator for his own machine.

TECHNIQUE FOR RAMUS OF THE MANDIBLE (Figs. 13, 14, 15, 16)


Fig. 13—Head and Cassette Position—Patient seated sidewise in chair, cassette resting on head-rest at angle of 45° with the floor and steadied with patient's hand on lower corner. Patient's head is thrown backward with the external ear flat against cassette. An imaginary line drawn through the two external auditory meatuses should be perpendicular to the cassette.

Figs. 14 and 15—Position and Direction of Tube—The tube is positioned under, and slightly anterior to angle of the jaw, so that the central ray will hit the cassette at a mesio-distal angle of 90° and vertico-horizontal angle of about 60°. The central ray will pass directly through the center of the ramus.

Fig. 16—Roentgenographic Anatomy—A clear view is obtained of the ramus of the mandible, the head of the condyle, and the glenoid fossa. The body of the mandible of the opposite side is superimposed on all the teeth except the two lower molars. The radiolucent area across the neck of the condyle is caused by the pharynx and is apt to be misinterpreted as a fracture in this region if one is not thoroughly familiar with his roentgenographic anatomy. The zygomatic arch can be seen running from the head of the condyle to the upper third molar. The anterior part of the glenoid fossa can be seen running down to the zygomatic arch. The radiopaque streak extending from the anterior surface of the ramus down across the two molar roots is caused by the internal oblique ridge.

This view of the mandible is indicated in cases suggestive of fracture involving the coronoid processes or head of the condyle, fracture of the zygomatic arch and any disturbances of the temporomandibular articulation.

It is also valuable in cases of osteomyelitis which attacks the ramus of the mandible more often than it does the body.



ADMINISTRATION OF ETHYL CHLORIDE ANESTHESIA

JULIUS A. GREENHOUSE, D.D.S.

Rochester, New York

ETHYL chloride is easily administered; its action is rapid, and there is ample time between the administration of the anesthetic and the return of sensation to the patient for the operator to perform whatever surgery is necessary.

CONTRA-INDICATIONS

Ethyl chloride was administered to children who were known to have heart disorders with no untoward results. The following conditions contra-indicate the use of ethyl chloride: (1) abnormally enlarged tonsils; (2) obstruction of air passages;

Stages in Complete Anesthesia—There are three distinct stages in complete anesthesia:

1. Stage of induction (analgesia):
 - (a) Period of excitement.
 - (b) Period of rigidity.
 - (c) Period of relaxation.
2. Stage of maintenance (anesthesia):
 - (a) First plane of surgical anesthesia: (1) eye reflex abolished; (2) oscillation of eyeball; (3) rhythmical breathing.
 - (b) Second plane of surgical anesthesia: eyeball fixed off center.

through the stage of induction, as far as the period of rigidity. The patient then returns to the stage of recovery, there being no period of relaxation nor stage of maintenance.

TECHNIQUE OF ADMINISTERING ETHYL CHLORIDE

Ethyl chloride is administered in small doses with large amounts of air.

Broadly speaking, there are two methods of administering ethyl chloride: what may be called the Jacob's or gauze technique and the Feldman or ether mask technique. Each method



Fig. 1—Patient in a semireclining position with body and head in a straight line.



Fig. 2—Middle finger placed on prop so as to prevent patient from pushing it out of mouth; at the same time little and third fingers rest under chin.




Fig. 3—Head of patient encircled in gentle grip between operator's body and left forearm.

- (3) diseases of the lungs; (4) diseases of the circulatory system, and (5) *status lymphaticus*.

CLASSIFICATION

Anesthetics are divided into two broad groups: general and local. By *general anesthesia* is meant the narcosis of the central nervous system. *Local anesthesia* affects a localized area by direct action of the agent on the terminal nerve endings. Ethyl chloride is being used both as a general and as a local anesthetic.

Classes of General Anesthesia—Flagg subdivides general anesthesia into two classes: complete and incomplete. Ethyl chloride is useful in dentistry because it can be used as an incomplete anesthetic.

- (c) Third plane of surgical anesthesia: eyeball fixed in central position.

3. Stage of recovery.

These stages and periods of anesthesia are not definite and complete, but blend into one another and there is considerable overlapping. The anesthetist must be constantly on guard.

Ethyl chloride is not desirable as a complete anesthetic. The margin of safety between the amount of the drug necessary to give complete relaxation and the amount producing toxicity is too narrow. Large doses of ethyl chloride rapidly given may cause cessation of respiration and spasm of the diaphragm.

Incomplete Anesthesia—In incomplete anesthesia, the patient is carried

has its advantages, but the mask technique will be discussed here.

1. The patient is seated in the chair and the chair tipped backward so that the patient is in a semireclining position with body and head in a straight line (Fig. 1). The head should not be too far backward or forward.

2. A mouth prop is placed in position on the occlusal surfaces of the teeth on the opposite side from the field of operation. The patient is instructed to bite down on the prop.

3. The middle finger of the left hand is placed on the prop so as to prevent the patient from pushing the prop out of the mouth; at the same time the little and third fingers rest under the chin (Fig. 2).

4. An ether mask (carrying from four to six layers of gauze) is placed in position on the middle finger and bridge of the nose, and the mask is held with the thumb at the upper end and the index finger at the base.

5. While the mask is being adjusted, the operator encircles the head of the patient so that it is in a gentle yet firm grip between the operator's body and left forearm (Fig. 3).

6. The ethyl chloride tube is held in the right hand and the anesthetic is either dropped or sprayed. When spraying the drug on the mask or gauze, the palm of the hand rests intermittently for a second or two, on the gauze, so that the heat of the hand prevents the ethyl chloride from freezing on the mask and shutting off the air or preventing the patient from getting the proper amount of anesthetic (Fig. 4).

7. When the anesthetic has been administered a mouth pack is in-

serted to keep the blood or tooth particles from being aspirated. The dentist then has about two minutes in which he can accomplish his operation (Fig. 5).

CLINICAL USE

1. Success in the administration of ethyl chloride depends on experience.

2. The chief usefulness of ethyl chloride is in operations of short duration (from 1 to 2 minutes), especially in children in whom muscular relaxation is not required.

3. As the drug acts rapidly, everything should be in readiness before induction is begun.

4. As the ethyl chloride is dropped or sprayed, consciousness is usually lost in from 30 seconds to 1 minute. When eyeballs are rolled up or down and a *slight snore* is heard (Fig. 6), the patient is ready for the operation.

5. Ethyl chloride is a cumulative anesthetic and should not be pushed too enthusiastically.

6. Usually about 3 cc. of the drug is administered.

ADVANTAGES

The following are the advantages in the use of ethyl chloride:

1. Safe.
2. No after-effects.
3. Rapid recovery.
4. Rapid action.
5. Ease of administration.
6. Small amount of anesthetizing agent necessary for the production of incomplete anesthesia.
7. Inexpensive.
8. Nonirritating.
9. Symptoms of anesthesia readily recognized.

DANGER

Danger in the use of ethyl chloride is due chiefly to interference with respiration or circulation. Cyanosis is not due to toxic effects from ethyl chloride, but to the incorrect position of the head, insufficient air or me-

(Continued on page 271)



Fig. 4—Ethyl chloride tube held in right hand and anesthetic being sprayed on mask. Palm of hand resting a moment on gauze to prevent ethyl chloride from freezing on mask.




Fig. 5—Insertion of mouth pack to keep blood or tooth particles from being aspirated.



Fig. 6—When eyeballs are rolled up or down, patient is ready for operation.



Fig. 7—Should cyanosis develop, mask is removed with forefinger of right hand and posterior third of tongue is moved forward to open throat and permit air to enter.



Figs. 8 and 9—Treatment demonstrated in the event of labored breathing.

"THE FIRST HUNDRED YEARS..."*

JOSEPH J. TOLAN, D.D.S., M.S.
Milwaukee



FIG. 1, A,
AGE, 2 DAYS




FIG. 1, B,
AGE, 12 DAYS




FIG. 2, A,
AGE, 3 MONTHS



FIG. 2, B,
AGE, 3 1/2 MONTHS



FIG. 3 A
AGE, 9 MONTHS




FIG. 3, B
AGE, 12 MONTHS




FIG. 1, C.
AGE, 6 WEEKS




FIG. 1, D
AGE, 2 MONTHS



FIG. 2, C.
AGE, 4 MONTHS



FIG. 2, D
AGE, 6 MONTHS



FIG. 3, C
AGE, 16 MONTHS



FIG. 3, D
AGE, 20 MONTHS

*This series of roentgenograms had its beginning more as a hobby than as a scientific venture. When the characteristic as well as unusual features began to unfold themselves, a more concentrated effort was made to obtain "types." There has been little chance to standardize. No attempt has been made to show normal tooth eruption and jaw development at stated ages. These roentgenograms were selected from a large collection because of individual, interesting factors; such as, the result of too early extraction of the deciduous molars and the contrasting picture of retaining the deciduous teeth too long. To "see" the crowns of the teeth calcifying and the teeth erupting; to realize that the tooth bud of the first permanent molar was in evidence at birth—these

and a dozen other observations, prompted the publishing of this series of roentgenograms showing the mouth as it may appear at each of the various age groups.

Fig. 1—Calcification of the cusps of the deciduous teeth and the development of the tooth bud of the first permanent molar.

Fig. 2—Completion of the development noted in Fig. 1. D shows the calcification of the cusps of the permanent first molar.

Fig. 3—Normal eruption of the deciduous teeth from 9 to 20 months



FIG. 4, A
AGE, 2 YEARS




FIG. 4, B
AGE, 30 MONTHS




FIG. 5, A
AGE, 3 YEARS



FIG. 5, B
AGE, 3½ YEARS




FIG. 6, A
AGE, 5 YEARS




FIG. 6, B
AGE, 5½ YEARS

Fig. 4—Compare B, appearance at 30 months, with A, C, and D. B demonstrates a late development and eruption of the teeth and a tardy development of the jaw itself. Compare the obtuse angle of the jaw with the normal angle in A (appearance at 2 years).



FIG. 4, C
AGE, 31 MONTHS




FIG. 4, D
AGE, 3 YEARS



FIG. 5, C
AGE, 4 YEARS




FIG. 5, D
AGE, 4½ YEARS



FIG. 6, C
AGE, 5 YEARS 9 MONTHS




FIG. 6, D
AGE, 6 YEARS

Fig. 5—This group demonstrates the calcification of the bicuspids.
A, Tooth bud of the second permanent molar. C, A good "normal."

Fig. 6—Progressive calcification of the bicuspids and second permanent molars.

RESTORING THE UPPER FIRST BICUSPID

JEROME M. SCHWEITZER, D.D.S.

New York




Fig. 1

IN RESTORING the upper first bicuspid when the tooth in position on each side is vital, the usual procedure is to replace the tooth by means of a fixed bridge. In this procedure the pontic is soldered to a mesio-occluso-distal inlay, or a mesio-occlusal inlay, or a three-quarter crown, which is placed on or in the second bicuspid, with a lug resting on an unprepared cuspid or in an inlay prepared in the cuspid with a




Fig. 2

Fig. 1—Additional retention of bicuspid inlay.

Fig. 2—Cuspid preparation.


Fig. 3—In the lingual view of the cuspid are seen two countersink holes with the finishing groove just below the incisal edge. The first bicuspid was extracted one week previous to taking the photograph. The bicuspid preparation with the mesial lock can be seen.

Fig. 4—Wax is pressed into position on the cuspid and trimmed as carefully as possible. A strip of rubber dam is drawn taut around the tooth. A hot instrument causes the wax to flow and better adapt itself to the tooth. The rubber dam is removed and two plain pins, which were fitted into the holes, are heated and pressed into the two holes and allowed to cool. The pattern is withdrawn.

Fig. 5—The entire lingual surface is then invested for casting. The investment is brought around to include everything but the pins and the surrounding wax of the lingual surface. When the investment is hard the pins are heated and pulled out and the holes are waxed.

Fig. 6—A single sprue pin is then inserted and the remainder of the wax is invested and cast.

Fig. 7—The bite is adjusted. Sufficient gold is left on the cuspid at the incisal to allow for trimming when the bridge is finished. An impression is now taken, a check-bite, and the bridge is constructed as indicated in the text.







Fig. 8—The different portions of the bridge taken apart. It is always wise to grind in an extra tooth in case one breaks. Note the lock lug. The bicuspid inlays receive this. Note the pontic attached to the cuspid inlay.

Fig. 9—The bridge cemented in the mouth. The incisal gold of the cuspid inlay has been trimmed and finished. The bite has been adjusted. Also, note that the bucco-lingual diameter of the pontic is smaller than the original bicuspid. Less trauma takes place when grinding the teeth. The porcelain pontic has been put in with only a small amount of gutta-percha. As this socket is not completely sealed, it is not wise to cement the tooth permanently. In a short time the tooth is removed to observe the socket. When the operator is satisfied that the healing is complete, then the tooth may be permanently cemented. Sometimes, porcelain must be taken away from the pontic as the socket heals; at other times, porcelain must be added to the pontic and reglazed, before the final cementing.

Fig. 10—An example of the same type of bridge on the right side.



lock lug on the distal of the cuspid. The inlay in the bicuspid must be retentive to hold the bridge, and the cuspid must have a considerable portion of the distal taken from it in order to allow for the lug. If no preparation at all is used on the cuspid, then the fear of decay beneath the lug which rests on it is always present.

I use a method at present which I feel overcomes these difficulties. If the adjoining teeth are in good condition, a mesio-occlusal inlay is prepared in the second bicuspid. This inlay should be made retentive by means of a lock in the mesial surface, or a countersink pin in the distal surface or mesio-cervical floor, or both, as security against loosen-

ing when the bridge is cemented (Fig. 1).

PREPARATION OF CUSPID

1. If the missing bicuspid is an upper left, the lingual of the upper left cuspid is ground away to allow for a thickness of gold; the mesial marginal surface and the incisal edge are not disturbed at all.

2. A groove is made, just distal to the mesial margin on the lingual surface, running parallel to the mesial margin, and two pins are countersunk, one near the incisal margin and the other near the cervical margin. The two countersunk holes must be parallel to the same plane in order to withdraw the pattern successfully.

3. The distal marginal ridge is removed with a stone, and a groove is placed here with a Tinker 71 bur,

running parallel with a line running labially from cervical edge to incisal edge.

4. In order to prevent showing gold at the incisal edge a finishing line is cut from the mesial hole to the distal groove and this line is below the incisal edge (Fig. 2).

5. A Steele's porcelain pontic is then ground in and soldered to the cuspid inlay with a lock-in lug in the bicuspid inlay. By reversing the order in this way the desired results are accomplished; namely: (1) the esthetic result is excellent; (2) the cuspid need not be cut deeply on the distal surface to allow for the lug; and (3) the bicuspid does not need extensive preparation, because it only supports a lock-in lug and not the pontic.

730 Fifth Avenue.

ADMINISTRATION OF ETHYL CHLORIDE ANESTHESIA

(Continued from page 265)

chanical blocking of the throat by the tongue or aspirated blood.

WARNING

1. Not all patients will react in the same manner to ethyl chloride. Alcoholic addicts are poor subjects. It is my experience that red haired patients have to be watched more closely than blond patients and blonds more than brunettes. Red-heads and blonds will offer a greater resistance to the anesthetic than brunettes.

2. Another observation is that not all ethyl chloride preparations produced by manufacturers are alike. I have not been able to ascertain whether this is due to lack of purity of the product, or age of the drug. One must find a preparation of ethyl chloride that is satisfactory and use it consistently for best results.

3. Ethyl chloride should be kept in a dry, dark place, away from heat or moisture.

14 Franklin Street.

EMERGENCIES

1. While administering an ethyl chloride anesthetic, should anything take place that would make the operator dubious as to the normalcy of the anesthetic, remove the mask at once.

2. Should cyanosis develop, remove the mask and with the forefinger of the right hand, move the posterior third of the tongue forward to open the throat and permit air to enter it (Fig. 7).

3. Should the operator notice difficult or labored breathing, the anesthetic should be stopped at once, and with the left hand (Fig. 8) the body of the patient should be pushed forward, so that the head is in position between the knees; at the same time the region of the diaphragm is massaged upward with the right hand (Fig. 9).

4. If, for any reason, the anesthetic has to be discontinued before the operation is begun or finished, do not continue with ethyl chloride. Either postpone the operation or use some other anesthetizing agent.


CONCLUSION

I have personally given about 3,500 ethyl chloride general anesthetics to patients ranging from 9 months to 15 years of age. Doctor Jacobs of the Forsyth Clinic in Boston reports about 195,000. Doctor M. Hillel Feldman has given in his private practice and in the Clinic of Lincoln Hospital, New York, more than 60,000 anesthetics of ethyl chloride and at no time was any difficulty encountered. Leeds Dental Hospital, England, has used ethyl chloride for fifteen years without a mishap.

CAST INDIVIDUAL IMPRESSION TRAYS

LEONARD S. FLETCHER, D.D.S.


Castle Shannon, Pennsylvania



MANY failures in prosthodontia are due to insufficient retention in the dentures, caused by the distortion of the impression in removing them from the ridges. For instance: When an upper impression adheres tightly to the tissue, we place the index finger

on the peripheral rim of the impression in the tuberosity region and try to force this portion of the rim outward and downward. Unless the tray or impression material is strong enough to withstand this pull, the impression will be distorted on removal. This will result in a distorted cast,

Legends on
opposite page



Technique

FIG. 1

Step 1. Take a snap impression of the ridge (A).

Step 2. Box-in snap impression with metal rim (B).

Step 3. Pour snap impression with plaster (C).

Step 4. Separate cast from snap impression (D).

FIG. 2

Step 5. Adapt one thickness of baseplate wax over the snap cast and dust freely with talcum powder (A).

Step 6. Adapt the second thickness of baseplate wax over the first powdered wax (B).

Step 7. Place the second piece of adapted wax (B) on the workbench with the tissue surface downward; box-in with a rim of soft metal, leaving at least one-fourth inch clearance between rim of wax and metal form (C).

FIG. 3

Step 8. Mix flasking plaster and pour over the wax within the metal form so as to have at least one-fourth inch of plaster at the thinnest area (A).

Step 9. Invert this half-invested wax form and remove any plaster that has run over the periphery onto the tissue surface. Allow this plaster containing the wax form to drop

to the bottom of the metal form. Paint the plaster with a good separating medium (B).

Step 10. Mix flasking plaster and pour over the wax to the height of the metal rim. The wax form is now full flasked and the metal rim removed (C).

FIG. 4

Step 11. Force a knife blade between the two halves of the mold at the center of the heel and separate (A).

Step 12. Remove the wax and cut a hole through the plaster rim at the distal of each tuberosity of the half of the mold from which the wax has been removed. When the metal is poured in one hole, the other will act as an air escape (B).

FIG. 5

Step 13. Close the two halves of the mold; hold together with a C clamp and stand it on the workbench (A).

Step 14. Melt the metal in a ladle and pour in the hole on one side of the mold until it overflows from the hole in the opposite side (B).

Step 15. Separate the two halves of the mold as mentioned in step 11 and break the plaster away from the cast tray (C).

FIG. 6

A. Rough cast tray.
B. Cast tray with excess metal ground away.
C. Original snap cast.

FIG. 7

The technique for making a cast individual tray for an immediate denture varies only in steps 5 and 6. The substitute steps are herewith given:

Substitute Step 5. From the snap impression (A) make a snap cast (B). Over this snap cast, adapt one thickness of Kerr's Utility Wax or Dentsply Boxing Wax and dust freely with talcum powder (C).

Substitute Step 6. Over this black wax adapt one thickness of baseplate wax (D).

FIG. 8

A. A snap cast of an immediate case.
B. A cast individual tray for same case.

FIG. 9

A and B. After adapting the second piece of baseplate wax, add an extension of wax from the base of the labial rim which will later form the handle of the tray when making the casting.

C. When making a cast tray with a handle make the sprue hole at the end of the wax handle extension, instead of at the heel of the mold.

FIG. 10

A. A desirable ladle in which to melt the tray metal.

B. An ingot of low-fusing special tray metal from which nine to twelve separate trays may be cast before remelting any of the trays.

and in the end, an incorrect adaptation of the denture to the tissue in the area of the postdam at that tuberosity.

This error can be eliminated with the use of a cast individual tray of sufficient firmness to prevent any distortion of the impression while dis-

lodging, regardless of the impression material used for registering the negative of the tissue.

SUGGESTION

It is preferable to have the plaster mold dry when pouring the metal. Pour the metal as rapidly as possible. After separating the mold into which

the metal has been poured, we may find that the tray did not fully cast. Should this occur, stand the half of the mold containing the partly cast tray on the workbench and melt the metal from the mold with the flame of a blowtorch. Close the mold and hold together with the C clamp, and make another pour.

STONE MODELS FOR INDIRECT CASTINGS

ROBERT S. HINES, D.D.S.

Cleveland, Tennessee

I WILL not try to settle the question of which die material is the better or more accurate, stone or metal. Each has its advantages and disadvantages.

Stone dies, when properly made, are to my mind by far more practical and are easier to make. They can be used for almost any type gold restoration, inlays, three-quarter and full crowns. Less time is required in making them, and they may be used immediately after they set—usually in fifteen minutes. Stone changes to some extent on drying; it is best, therefore, that the pattern be carved before the die thoroughly dries; that is, from six to eight hours after it is made.

If stone is used as a die material, inlay wax may be used as an impression material. After the stone has set sufficiently, the band is slit and the pattern carved; the impression wax is used for carving. By using inlay wax for taking the impression, one is able to obtain a finer, more accurate impression than with any other material.

TECHNIQUE

1. A clean-cut impression is obtained in inlay wax.

2. Thoroughly dry with blasts from a chip blower.

3. A thin coating of light machine oil is then painted on the impression with a fine camel's hair brush.

4. A simple vibrator for use in the handpiece may be made in the following way:

A. Flatten an orangewood porte-polisher (Fig. 1).

B. Insert the orangewood porte-polisher in a mandrel (Fig. 2).

C. Place the inlay wax, copper band impression in the tin lid from a small box.

D. Mix the artificial stone into a stiff mix on a cement slab.

Heartsill Building.



Fig. 1




Fig. 2




Fig. 3




Fig. 4

Fig. 1—Orangewood porte-polisher.

Fig. 2—Porte-polishing mandrel.

Fig. 3—Vibrator assembled, showing handpiece also.

Fig. 4—Copper band impression on box lid.

E. Using a beaver tail burnisher, place small amounts of the stone in the impression and vibrate to place by striking the tin upon which the impression is placed with the vibrator (Fig. 3).

5. A piece of stone is rolled be-

tween the fingers into a handle; this is placed on top of the full impression.

6. After the stone has set, the copper band is slit and the die is separated from the impression.

7. The die is oiled thoroughly; the wax impression replaced, and the pattern carved.

The Editor's Page

GLOSSODYNIA (hyperesthesia, burning tongue, or glossopyrosis) is a condition that is sometimes encountered in dental practice. A burning or painful sensation of the mucous membrane over the palate or ridges is an occasional complaint of persons wearing full dentures. The hyperesthesia in the edentulous is thought by some to be produced by pressure of the denture base on the nerves and vessels that emerge through the anterior and posterior palatine foramina. Burning sensation under denture bases has been frequently explained on the basis of low tissue tolerance to mechanical and possible chemical irritants. Persons whose "threshold for pain" is low and whose nervous irritability is high are the most common sufferers from glossodynia or hyperesthesia of the oral mucous membranes.

All observers agree that the majority of persons afflicted are women. Women at or beyond the menopause¹ are more often sufferers than younger women. Blair² believes that the condition is encountered in anemic, poorly nourished, neurotic women. In Gilpin's³ series of forty-eight patients from the Mayo Clinic, thirty-seven were women and the average age of the whole group was 54.3 years.

Both Prinz and Greenbaum⁴ and Gilpin⁵ comment on the relationship between burning, painful tongue and gastric hyperacidity. Gilpin quotes Sellei⁶ who postulated that glossodynia represents a reflex disturbance from the stomach over the vagus nerve to its nucleus in the fourth ventricle and then to the nucleus of the fifth cranial nerve and over the branches of the trigeminal nerve to the tongue. Sellei found, and his observations, according to Gilpin, were confirmed by Morelli,⁶ that the administration of alkalis and dietary therapy relieved the burning sensation.

Prinz's description of the typical symptoms is excellent:

¹Osler, William; and McCrae, Thomas: *The Principles and Practice of Medicine*, Ninth Revised Edition, New York, D. Appleton & Co., 1920, page 453.

²Blair, V. P.: *Surgery and Diseases of the Mouth and Jaws*, Third Edition, St. Louis, C. V. Mosby Company, 1918, page 533.

³Gilpin, S. F.: *Glossodynia*, J. A. M. A. 106:1722 (May 16) 1936.

⁴Prinz, Hermann; and Greenbaum, S. S.: *Diseases of the Mouth and Their Treatment*, Philadelphia, Lea & Febiger, 1935, page 458.

⁵Sellei, Josef: *Durch Hyperazidität verursachtes Zungenbrennen*, Deutsche med. Wchnschr. 2:1758 (October 19) 1928.

⁶Morelli, Gustav: *Reflectorische Ueberempfindlichkeit der Zunge (Hyperaesthesia reflectoria linguae)*, abstr. Zentralbl. f. Haut.-u. Geschlechtskr. 29:309, 1929.

The outstanding symptom in all cases is typically characteristic, i.e., a burning sensation within the anterior third or the tip of the tongue and near its borders. Occasionally, the painful region extends to the soft and hard palate and to the vermillion border of the lower lip. The diffuse pain is usually described as being of a burning character, sometimes as itching or boring and, at other times, as presenting all variations of an indefinite diffuse painful sensation. The pain is more or less constant but never paroxysmal. The patient gives the impression of being deeply depressed. There are no points douloureux to be observed. The pain usually disappears during animated conversation or when the patient's attention is concentrated upon some inspiring subject in which he is vividly interested. Paresthesia is frequently present.

A common complaint is dryness of the mouth resulting from more or less inhibition of the flow of saliva, while taste and thermal sensibility are not altered by the disease. On the other hand, the pain is usually increased when the tongue is brought in contact with liquid or solid food.

Gilpin mentions the empirical treatment that patients with this condition have frequently suffered at the hands of dentists and physicians. Persons with a burning sensation of the tongue, palate, or oral mucous membranes frequently believe that some dental condition is responsible: a crown; an edentulous space; a bridge. Dentists have been known to recommend radical treatment in cases of "heavy dentistry" on the basis of a possible electric phenomenon produced by dissimilar metals in restorations. Physicians may recommend extractions "on general principles."

If burning sensations in the mouth are of psychoneurotic origin (Gilpin) or represent a reflex disturbance or are produced by a general metabolic derangement, it is evident that local dental treatment should be extremely conservative. To extract teeth and subsequently insert extensive restorations in a mouth already in discomfort, and thus introduce an additional mechanical irritant is not sane treatment. All authorities agree that local treatment is futile. Glossodynia is another example in which the combined skill of the dentist and physician is required.

A REBASING TECHNIQUE

P. N. OGLE, D.D.S.

Knoxville, Tennessee




Fig. 1

Fig. 1—Palatal view of full upper denture prepared for rebasing technique. The palate has been cut away to give relief over the anterior palatine and posterior palatine foramina and over the bony eminence of the hard palate. Metal plate waxed in position.




Fig. 2

Fig. 2—Lingual aspect of full upper denture prepared for rebasing. The metal on the palatal surface is held in place with sticky wax. The holes in the metal plate allow for escapement of plaster when the impression is taken.

REBASING full upper dentures is a service that every dentist is frequently called on to perform; without having a definite technique to follow, it is difficult to obtain good results. Papers have been written and many clinics have been given on full upper impression techniques, but to my knowledge no definite plan of procedure for rebasing has ever been given. A simple technique that I have been using for some time has proved satisfactory. This technique can be used in taking an impression for a full upper denture as well as for rebasing.

TECHNIQUE

1. The part of the palate that covers the hard area is cut out, and the opening is extended to include the anterior and posterior palatine foramina (Fig. 1).

2. A piece of metal (tin or aluminum), larger than the opening in the palate is cut to conform to the under side of the denture and fastened with sticky wax.

3. Holes are bored in the metal at the foramina and hard palate to allow for the escape of the excess plaster while taking the impression (Fig. 2).

4. The patient is instructed to hold the denture in place with pressure until the plaster sets. This will place

Medical Arts Building.

Fig. 3—Modeling compound impression relieved in the shape of a "T" to include bony eminence on the hard palate and the principal nutrient foramina. In this technique the holes are drilled through the metal impression tray to allow for the escapement of plaster on taking the "wash" impression.




Fig. 3

the stress only on the stress-bearing area, and will not allow pressure on the hard palate or over the foramen; but it will give adaptation to soft tissues.

5. The impression or wash is then taken in the usual way.

IMPRESSION FOR FULL UPPER DENTURE

Modeling compound is used for

the initial impression. The compound is chilled, trimmed, and cut out for relief in the same shape as described in the rebasing technique. Holes are then bored in the metal impression tray to allow the escape of excess plaster. Trenches should not be cut at the postdam as the perforations in the tray will take care of the excess (Fig. 3).

RELIEF OF PAIN



The Dentist's
NO. 2 FUNCTION*

ACETANILID

Promptly Relieves Pain Of All Nerve Origin

*Fantus holds that the relief of pain is second only to the saving of life. (J. A. M. A. Dec. 22, 1934.) The dentist, when choosing an analgesic, considers these major factors—efficiency—promptness—safety—non-habit-forming properties.

Acetanilid fully meets these requirements of an analgesic. It promptly relieves all pain of nerve origin by blocking the synapses and preventing pain sensation from reaching the higher centres. It does this safely, as widely cited in the literature, without injury to heart, kidneys, brain centres, blood pressure, etc. It has never been proven to be habit-forming.

For rapid diminution of pain in a safe, economical manner, the dentist has at his disposal a scientifically produced pharmaceutical. Acetanilid, with its synergists, caffeine and bromides, for maximum results with minimum dosage, is given in a palatable, refreshing drink. Citrates bring about favorable alkalization.


Generous trial supply is at your disposal upon request.

(Please use coupon on page 286)

EMERSON DRUG CO., Baltimore, Md.

PHOTO COURTESY UNITED
AIRCRAFT CORPORATION

Men stake
their lives
on this
engine



—made finer by "Specific" alloys!

In this Pratt & Whitney 1000 h.p. "Twin Wasp" are 19 casting alloys in the aluminum group alone, as opposed to 2 used in airplane engines during the World War. The reason? . . . modern metallurgy has progressed so far that it is no longer necessary to tolerate alloys that are only "good enough" for a certain use.

The same thing holds true of modern dental alloys. For example, the Ney-Oro "B" golds listed below enable you to make better $\frac{3}{4}$ crowns, dummies, pontics, bridge abutments, etc., because they suit the requirements of such work exactly. They have just the ductility, strength and hardness necessary for the finest results in these particular types of restorations.

For $\frac{3}{4}$ Crowns, etc.

NEY-ORO B	\$2.30 DWT.
NEY-ORO B-2	2.05 DWT.
NEY-ORO B-20	1.71 DWT.

NEY'S *Specific* ALLOYS

EACH DESIGNED TO MEET A *Specific* REQUIREMENT *Exactly*



Wherever You'd Use a Plaster Wash

Dr. Kelly's Impression Paste gives you an exact registration of details of the soft tissues; helps you to avoid most of the grief of denture work, and to turn out a superior result every time. The NEW ready-mixed Paste in tube form makes setting speed easy to control; causes no smarting; stays soft on the slab for long periods; hardens quickly in the mouth. Users are most enthusiastic. At dealers, \$2.50. Try it; we guarantee your satisfaction. Kelly-Burroughs Laboratory, Inc., 143 N. Wabash Ave., Chicago, Ill.

USE "KELLY'S PASTE"

ABOUT OUR CONTRIBUTORS

GEORGE W. MATTHEWS, D.D.S. (Northwestern University, 1927) was the co-author of an article, ORAL SURGERY: REPORT OF CASES, with Harry Bradford, D.D.S., which appeared in the May, 1936 issue, and his professional biography was given in that number of this magazine.

JULIUS A. GREENHOUSE, D.D.S. (College of Dentistry, University of Iowa, 1923) interned at the Rochester Dental Dispensary and after nine years of general practice again interned at Lincoln Hospital, New York City (1933-1934). Doctor Greenhouse is a member of the A. D. A.; the American Society for the Advancement of General Anesthesia in Dentistry; County of Monroe Dental Society; and the Rochester Dental Study Club where he is the group leader of the section on anesthesia and surgery. Doctor Greenhouse's practice stresses dental anesthesia and surgery.


JOSEPH JAMES TOLAN, D.D.S. (University of Michigan, 1923; M.S., 1925) is a member of the American Dental Association and the Wisconsin Academy of Dentists. Doctor Tolani spent three years in the department of oral surgery at the University of Michigan Hospital under Doctor Chalmers J. Lyons. Doctor Tolani now specializes in oral surgery and exodontia.

JEROME M. SCHWEITZER, D.D.S. (New York University College of Dentistry, 1918; B.S., College of the City of New York, 1927) contributed an article on EXTENSIVE PROSTHETIC RESTORATIONS: RAISING THE BITE in the April, 1936 issue of the DIGEST. Doctor Schweitzer's professional biography appeared in that month.

LEONARD S. FLETCHER, D.D.S. (University of Pittsburgh, 1924) is a member of the American Dental Association and the Odontological Society of Western Pennsylvania, the Pennsylvania State Society, and the South Hills Dental Society. Doctor Fletcher has given postgraduate courses at the University of Pittsburgh as well as private courses on immediate denture construction. He has previously contributed to this magazine (August, 1933) and to other periodicals.

ROBERT S. HINES received his D.D.S. in 1935 at the University of Tennessee, College of Dentistry. Doctor Hines is a member of the American Dental Association and has a general practice.

P. N. OGLE received his D.D.S. in 1923 at the Louisville College of Dentistry. Doctor Ogle is a member of the A. D. A. and component societies, and is on the staff of Fort Sanders, Knoxville General, and St. Mary's Hospitals. Doctor Ogle practices general dentistry.



COPY FREE ON REQUEST

Dependable sources of Vitamin D effect under the Steenbock Irradiation Process include:

Viosterol and Viosterol products produced under the Steenbock patents by Abbott, Mead Johnson, Parke-Davis, Squibb and Winthrop.

Irradiated Evaporated Milk produced by Borden, Carnation, Nestle, Pet and Wilson.

Irradiated Vitamin D fluid milk—now available in most large and many smaller cities.

Metabolized Vitamin D fluid milk available in nearly 300 cities.

Other Vitamin-D-enriched foods include: Coco-malt; Dryco powdered milk; Fleischmann's Irradiated Foil Yeast; Ovaltine; Quaker Farina, Muffets whole wheat biscuits, and Quaker and Mother's Rolled Oats; Sunfed Flour.

**W I S C O N S I N A L U M N I
R E S E A R C H F O U N D A T I O N ★**

* A corporation not for private profit . . . founded in 1925 . . . to accept and administer, voluntarily assigned patents and patentable scientific discoveries developed at the University of Wisconsin. By continuous biological assays, the public and professional confidence in accurately standardized Vitamin D is maintained. All net avails above operating costs are dedicated to scientific research.

(Please use coupon on page 286)



**A VITALLY
IMPORTANT
DEVELOPMENT
IN DENTISTRY**

—the use of Vitamin D in tooth and bone building, nourishment and protection—is authoritatively treated in this new booklet, made up of logically arranged abstracts from scientific papers by recognized authorities. This significant booklet is in broad demand throughout the profession; copy free to dentists and physicians upon request.

**M A D I S O N
W I S C O N S I N**



Are you a Good Dental Patient?

You are a dentist. You know infinitely more about the health value of good dentistry than any layman. Yet, do you willingly have your teeth cared for at regular intervals? Or are you subject to the same inhibitions which make dentistry an ordeal to the average lay patient?

The next time you go to a dentist select one of your colleagues who is equipped with a McKesson Euthesor. Contrast the treatment under gas analgesia with your previous experiences in a dental chair. Put yourself in the frame of mind of the average patient, whose attitude toward dentistry is greatly influenced by the degree of pain he suffers during dental treatment. Let your own experience tell you whether or not McKesson equipment would be a good investment for you.

One of the realities which any dental practice must face is the fact that *people do not want to be hurt*. The elimination of operative pain is a primary step in making dentistry more appealing to the public.

With McKesson analgesia patients are put at ease. Nervous tension is replaced by calm assurance. Patients' dread of dental treatment is eliminated and operating conditions are improved.



McKesson literature will tell you more about nitrous oxid-oxygen analgesia, its technique, possibilities and cost.

Return the coupon on page 286.

McKESSON APPLIANCE CO.
TOLEDO, OHIO

A. D. A. CONVENTION REPORTS

"DENTAL GROUPS WARMLY DEBATE POLICY CHANGES"

"DELEGATES PASS MEASURE FOR NEW EDUCATION COUNCIL SETUP"

"Two bitterly opposed factions threw the American Dental Association convention into an afternoon of acrimonious debate . . . over a proposal embracing far-reaching changes in dental education, foundation stone of the profession.

"Designed to elevate the standards of dental education by centralizing control in the association itself, the proposal was passed after a stormy session of the House of Delegates, legislative body of the association, at the Hotel St. Francis.

"It provides:

"1—For reduction of the Dental Educational Council to a membership of seven. The council was hitherto composed of nine members, three each from the American Dental Association, the National Association of Dental Examiners and the American Association of Dental Schools.

2—That all appointments to the council be made by the board of trustees of the American Dental Association. The appointive power has hitherto rested in the three organizations represented on the council.

"3—That the council include three representatives of the American Dental Association and two representatives each of the other two bodies.

"Since 1909, the council has had full control over the classification of dental schools throughout the country and has thus wielded great power in setting the standards for admission to the profession. Under the new setup, this power will vest in the dentists themselves.

"RECOMMEND CHANGE"


"The changes were recommended in a report from a reference committee on education, headed by Dr. T. E. Purcell, dean of the St. Louis University Dental School, and including Dr. William A. Davis of Lansing, Mich., and Dr. A. Y. Russell of Baltimore, Md.

"Strenuous opposition came from Dr. Albert L. Midgley, representative of the examiners' association on the council, who demanded equal representation for the three groups, and from Dr. Shepherd W. Foster of Atlanta, Ga., representing the school association on the council.

"Dr. Foster condemned the proposal and resigned from the council forthwith.

If you are not a subscriber
to
THE DENTAL DIGEST
you will find a convenient
order blank enclosed.
Note special offers.

*In Cases Requiring
Draining or Dressing*



CAMPHO-PHENIQUE

Antiseptic • Anodyne • Healing

is Second to None

After post-operative treatment, in gingival infections, pus sockets and other oral infections, you can rely on CAMPHO-PHENIQUE liquid to arrest infection, soothe the pain and stimulate tissue repair . . . without irritation.

Try CAMPHO-PHENIQUE, doctor . . . as a swab for bleeding surfaces . . . as a gauze packing after suture removal, as a dressing after cutting. Its efficacy will invite your steady use.

CAMPHO-PHENIQUE is available in 1, 2 and 4 ounce bottles at all Drug Stores and Dental Supply Houses.

Safeguard your work by adequate follow up recommendations for intervals between office treatments.

Campho-Phenique Company
500-502 North Second Street
St. Louis, Mo.



Knowledge Applied

THE VALUE OF KNOWLEDGE is best measured in its application to the common good.

The public, we believe, is aware of the major facts about the prevention of tooth decay. There remains, however, a need for something to stimulate them to a wider application of this knowledge.

To provide this needed stimulus, the House of Squibb has crystallized these major facts into a simple, workable plan. Inspired by the encouraging fact that Most Tooth Decay Can Be Prevented, this plan is daily being carried into millions of homes by our newspaper, magazine and radio advertising.

Naturally, any such plan must include the use of a dentifrice. Squibb has spent many years developing dentifrices that conform to the highest standards of the dental profession, and has now developed a tooth powder, containing the same essential ingredients as Squibb Dental Cream.

This is the Plan We Advocate

1. Go to your dentist and follow his advice. This will include the kind of toothbrush to use, and how to use it; what kind of dentifrice to use, and what kind not to use; and whether you should supplement your own home treatment with the use of dental floss and oral perborate.
2. Check your diet with your physician or dentist—to be sure your system is getting the elements essential to the health and strength of your teeth.
3. Brush your teeth thoroughly, at least twice a day, and be sure you use a dentifrice scientifically prepared to clean teeth effectively, and safely.

SQUIBB DENTAL CREAM · SQUIBB TOOTH POWDER

The Priceless Ingredient of Every Product is the Honor and Integrity of its Maker


*it's sound buying to
modernize with
a Castle...*

**... AND "GOOD PRACTICE"
TO INVEST IN *Cast Bronze*,
FULL-AUTOMATIC Value.**

Even though your old sterilizer may still boil water, just give it a critical "once over" and compare its appearance to a modern Castle recessed model. Picture the difference in *your* office... the reaction of patients... as well as your own sense of safety and pride.

And as to value, a Castle is lifetime. Its boiler is a solid Bronze Casting—you'll never have trouble from leaks and no up-keep cost... and the control is Full-Automatic, which means that it runs itself. No 3-heat switch, yet it does automatically all that a control switch should do if remembered.

Write for new book "Modern Sterilization" showing sterilizers for every budget. Wilmot Castle Company, Sterilizer Headquarters, 1109 University Ave., Rochester, N. Y.



CASTLE STERILIZERS

50 Years of Quality Leadership

(Please use coupon on page 286)

A FREE CAN

Lee Smith Foil Coat—Improved—is a great help in prosthetic work.

No more need for foiling models for vulcanite work; no more wrinkles from the use of cellophane. Valuable time saved.

You will want to try Lee Smith Foil Coat—Improved—so if you will buy a 3 oz. can of Foil Coat—Improved—and a 3 oz. can of Foil Coat Thinner, for the regular retail price of \$1.90, we will give you another dollar can of Foil Coat—Improved—free.

The offer is of necessity open for a limited time only.

Use coupon on page 287.

LEE S. SMITH & SON MANUFACTURING CO.
7325 Penn Avenue Pittsburgh, Pa.

"MOVE SUPPORTED"

"Dr. Marcus Ward of the University of Michigan, also representing the school group on the council, vigorously supported the reference committee's recommendations. Dr. Ward lauded the new arrangement as a great forward step for the profession and as a close pattern of the system used by the American Medical Association.

"Dr. Purcell said:

"This committee is simply attempting to give the dental profession the main voice in its own affairs. Control of dental education should rest mainly with this association and not with the dental schools or the dental examiners."

—From: *San Francisco Chronicle*, Thursday, July 16, 1936.

"CHIEF CONDEMNS COMPULSORY HEALTH INSURANCE"

"DELEGATES... GUARD PROFESSION"

"America's dentists, assembled in San Francisco for their seventy-eighth annual convention, swung into action yesterday bent on drafting a program that will both safeguard their profession and adequately provide for human needs.

"Keynotes on the program were sounded in the presidential message of Dr. George B. Winter of St. Louis, delivered before the convention's first general session in the Hotel St. Francis.

"Concentrating on dangers confronting the profession through possible legislative enactment, Dr. Winter uttered a sweeping condemnation of compulsory health insurance and called upon the delegates to prepare for combat against such measures.

"We are on the threshold of a period," he said, "in which the fate of dentistry and medicine is held in the balance. We must be ready to participate in the struggle and to properly guide the destiny of our profession.

"The association has compiled a wealth of material showing the harmful results of State medicine and health insurance."

"For successful resistance of movements toward socialized dentistry, Dr. Winter looked to a close alliance with the medical profession and a vigorous educational campaign.

"SOCIALIZED DENTISTRY"

"His words were echoed by Dr. Sanford Moose, president of the California State Dental Association.

"Referring to the appointment of State legislative committees to study

⁽¹⁾ The Principles of DENTAL MEDICINE

By F. W. BRODERICK, Hon. Dental Physician to the Royal Victoria and West Hants Hospital, London, England.

The new Second Edition covers all the advances in dental medicine and its relationship to general medicine. An especially fine discussion of the etiology, prevention and treatment of dental caries and pyorrhea is included. The book is divided into three parts, the first section covering General Physiological Considerations. Here the new material on Acidosis and Alkalosis, the Colloidal State, Endocrine and Vegetative System, and Calcium Metabolism is given. Part II is devoted to Dental Medicine. Part III is given over to the Relation of Dental to General Medicine. Critics agree that this is one of the most comprehensive books ever published. Truly it is a work deserving a place in every dentist's library.

575 pages, 27 illustrations. Price \$7.50.

Contents: Preface. Prologue. Part I—General Physiological Considerations. The Problem. Reaction. Acidosis and Alkalosis. The Colloidal State. The Endocrine System. The Vegetative System. Diathesis and Constitution. Calcium Metabolism. The Nature and Origin of Disease. Part II—Dental Medicine. Introduction. Etiology of Dental Caries. Etiology of Pyorrhea Alveolaris. A Summary of the Previous Four Chapters. The Method of Investigation. An Introduction to Chemotherapy. The Preventive Treatment of Dental Caries. The Treatment and Prevention of Pyorrhea Alveolaris. The Aim of Medical Treatment. Part III—The Relation of Dental to General Medicine. Civilization as a Factor in the Problem. Dental Sepsis as a Factor in the Etiology of Disease. The Teeth in Allergic Diseases, Arthritis, Tuberculosis. The Teeth in Malignant Disease. The Teeth in Diseases of the Alimentary System. The Teeth in Diseases of the Respiratory System. The Teeth in Diseases of the Nervous System. The Teeth in Diabetes. The teeth in Certain Diseases of the Skin. The Teeth in Diseases of the Eye. The Teeth in Acute Infectious Fevers. Some Illustrative Cases.

A MEMO ABOUT SOMETHING YOU HAVE BEEN PLANNING TO DO

Most of us have traveled a long way since the day we left college with our diploma tucked under our arm. Dentistry, too has travelled far in the intervening days. Day by day new discoveries have been made, new techniques developed. The cumulative result of this progress has brought many changes and advantages to practically every field of dental practice since the days when most dentists were learning the fundamentals in dental college.

The record of this progress is to be found in modern dental books. The help, instruction and counsel of the best minds in the profession are available to the forward looking dentist at little cost.

How many times have you decided that some particular book would be helpful to you—in some cases decided to order it, and then, for some reason or other put it off?

How many times have you planned to fill up the gaps in your library, only to postpone definite action?

We need not remind you of the value of practical up-to-date books. Nor can we tell you the books you need most. You know more about this than we do. We can, however, remind you that books are easy to put off. They are also easy to buy.

Easy Payment Plan

The Mosby Easy Payment Plan enables you to stock your dental library with the working tools that you need—and to do this immediately. This plan entails no financial hardship. If your order is \$15.00 or less the monthly payments need be only \$3.00. If your order calls for more than \$15.00 worth of books divide the entire amount by six to arrive at the monthly payments. No down payment is necessary. Simply fill in the attached coupon, listing the books you want, then write your name and address plainly. The books you select will be forwarded immediately—PREPAID.

(2) ANESTHESIA IN DENTAL SURGERY

By Sterling V. Mead, Washington, D. C.

There is no subject today of greater economic importance to the forward looking dentist than anesthesia. The selection of the anesthetic, the choice of technic, the accuracy of administration, all have a bearing on the results. In this book, recently off the press, is presented a complete manual on anesthesia in dentistry, both local and general. There is no anesthetic best for all cases, but there generally is an anesthetic best for each operation. This modern book on anesthesia provides the solutions to your daily anesthetic problems. Profusely illustrated, thorough in its coverage it is a book that will pay for itself many times over.

482 Pages Price, \$6.50 144 Illus.

(3) DISEASES OF THE MOUTH

By Sterling V. Mead, D.D.S., M.S., B.S., Washington, D. C.

The ability to recognize pathological disturbances and to diagnose and treat these conditions successfully is a fundamental necessity in modern dental practice. This book supplies the dentist with the fundamental principles and practical knowledge necessary for routine work in making a correct diagnosis of abnormalities within the mouth, thus permitting a correct prognosis so that proper treatment may be applied more promptly and more efficiently. Every phase of diagnosis and treatment of diseases of the soft tissue is gone into carefully.

322 Pages 40 Color Plates 523 Illus.
Price, \$10.00

(4) ORAL SURGERY

By Sterling V. Mead, Washington, D. C.

This book brings oral surgery well within the range of possibilities of the average dentist. Operations which were considered hazardous and beyond the province of the dentist a few years ago, are now readily performed. Day by day more oral surgery is being entrusted to the dentist. This book brings oral surgery well within the range of possibilities of the average dentist. It is more than a comprehensive presentation of the subject. It is a step-by-step guide. Every operation is described in its progressive steps. Having this guide post in oral surgery means opening additional avenues of work and income.

1087 Pages Price, \$12.50 403 Illus.

(5) PRACTICAL PEDODONTIA

By Floyd Eddy Hogeboom, Professor of Children's Dentistry, College of Dentistry, University of Southern California.

Children's dentistry is a phase of practice the general practitioner can ill afford to neglect. Children of today grow up and become the fathers and mothers of tomorrow. Many important advances have been made in this field of dentistry. The newer methods of handling juvenile cases are discussed in this book which thoroughly covers the subject, including deciduous teeth, cavity preparation, root canal work, endocrinology, and public health dentistry. 328 Pages, 249 illustrations. Price, \$6.50

(6) PRACTICAL ORTHODONTIA

Dewey & Anderson

New Fifth Edition Revised by George M. Anderson, D.D.S., Professor of Orthodontia, Baltimore College of Dental Surgery. With the assistance of the following contributors: B. W. Weinberger, Sidney Riesner, E. H. Broadbent, H. E. Kelsey, Rudolf Kronfeld, A. F. Rogers, Earl W. Swineheart, C. F. Wright, E. B. Arnold and E. A. Kittowski.

This book is wedded to no one idea or system, its purpose being to present in one volume authoritative information which will aid the general practitioner and the dental student in intelligently understanding the problems of orthodontia, and to assist the orthodontist in the interpretation and control of orthodontic cases. Starting with the first chapter, devoted to the principles of orthodontia, the subject is treated progressively through 25 chapters, thus giving the reader the etiology and principles involved before demonstrating the actual treatment of cases. Fixed and removable appliances are clearly described in both their construction and application. The treatment of cases is carried through step by step from beginning to end.

514 Pages Price, \$8.50 586 Illus.

(7) PROSTHETIC DENTISTRY

By Ira Goodsell Nichols, D.D.S., Champaign, Illinois

An encyclopedia of full and partial denture prosthesis. Forty-five of the leading prosthodontists in America and Europe have collaborated with Dr. Nichols in the preparation of this work. It gives the general practitioner, prosthetic dentist, and laboratory technician a definite and scientific procedure in the construction of artificial dentures, contains the best in methods, materials, and instruments advanced by leaders in the profession. The technic is explained step-by-step and is thoroughly illustrated. 688 pages, 830 illustrations, 3 color plates. Price, \$12.50.

THE C. V. MOSBY COMPANY, 3525 Pine Blvd., St. Louis, Mo.

DD-8-36

Gentlemen: Send me the following books: Nos. 1 (), 2 (), 3 (), 4 (), 5 (), 6 (), 7 (), charging my account. I agree to pay for these in accordance with your Monthly Payment Plan.

Dr. _____ Address _____

(Or please use coupon on page 287)

How Much Does Dental Cement Cost YOU? . . .



The answers to many other pertinent questions about the characteristics and use of crown and bridge cements are found in this booklet, said by many to be the most complete encyclopedia of its kind.

Do you know the method to use in determining your cost of cement per cementation?

It is not a difficult computation at all but one with which dentists generally are not familiar.

And it is important, for the difference between a profitable year and one of service without profit is often made up of little matters which considered individually seem "too small to bother with."


How to figure cementation costs is clearly shown in the booklet "Questions and Answers" issued by this company for the information and guidance of the profession. It may surprise you to learn that the "cheapest" cements are often the most expensive even when judged without any consideration of quality.

We will gladly send you a copy of "Questions and Answers" with our compliments for your office library. Simply tell us to whom and where to mail it. The W. V-B Ames Company, Fremont, Ohio.

(Please use coupon on page 287)

AMES DENTAL CEMENTS

the PUMICE
does the Work!



There is no substitute for fine Pumice as a prophylaxis material. It has every characteristic necessary for this work and when used in the form of Buffalo Prophylaxis Paste, it not only does a splendid cleaning job, but it leaves a pleasant remembrance in the mouth of the patient.

A generous quantity in a handsome opal glass jar for \$1.00 at your dental dealer.

Buffalo Dental Mfg. Co.

Kehr and Urban Sts.

Buffalo, N. Y.

Use coupons
on pages 286 - 287
when writing to
advertisers.

This service
is extended
to readers
at their request.

It enables you to
preserve your copies
of
THE
DENTAL DIGEST

the feasibility of socialized dentistry in California, Dr. Moose said:

"If any form of sick insurance is presented to the Legislature, the dental profession will fight to insist that it provide protection for the standards of dental health service now maintained in California. We will likewise insist on the preservation of the standards already fixed for those within our ranks.

"In short, we are determined never to be forced down to the standards of European countries which have adopted a health insurance or panel system."

"WAR ON CHARLATANS"

"Dr. [C. Raymond] Wells appealed to the convention for alertness and decisive action in driving 'quacks and charlatans' from the profession and for legislation which would assure the American dentist the right to practice without interference from outside agencies."

—From: *San Francisco Chronicle*, Wednesday, July 15, 1936.

"IMPROPER APPLICATION"

"Dr. Hartman blamed failure of the solution to produce complete anesthesia on improper application. It has been used effectively in all parts of the world, despite charges of the skeptical, he said."—From: *San Francisco Chronicle*, Tuesday, July 14, 1936.

"HERE'S WAY TO BRUSH TEETH"

"A man whose life work is to make people brush their teeth hard and often gave his ideas to a scientific section of the American Dental Association yesterday.

"Dr. Frederick A. Bricker of Los Angeles declared:

"Use as hard a tooth brush as you can buy.

"Work the bristles between the teeth, to force the blood out of the gums, thereby effecting circulation.

"Good circulation will prevent pyorrhea, the most common cause of losing teeth.

"Brush as vigorously as possible without pain or discomfort. To cause pain is to cause irritation and consequently harm.

"Remember, the gum tissue is as strong as that covering the palm of the hand and sole of the foot. Don't be afraid of a thorough massage."

—From: *The San Francisco Examiner*, Thursday, July 16, 1936.

"DENTISTS ELECT NEW OFFICERS"

"DOCTOR C. W. CAMALIER NAMED PRESIDENT FOR 1937-38 TERM"

"Dr. C. Willard Camalier of Washington, D. C., was elected president of the American Dental Association for the 1937-38 term . . . in the final business session of the association's seventy-eighth annual convention at the Hotel St. Francis.

"Dr. Raoul H. Blanquie of San Francisco was unanimously elected first vice president for the ensuing year.

"A private practitioner, Dr. Camalier was opposed only by Dr. Marcus L. Ward, former dean of the University of Michigan Dental School. His election was a reflection of the dentists' desire to fill key positions of the association with men engaged in private practice."

"COUNCIL REDUCED"

"The delegates also stamped final approval on a plan reducing the Dental Educational Council from nine to seven members by allowing the association three representatives on the Council, with two each from the American Association of Dental Schools and the National Association of Dental Examiners. The three groups were hitherto represented equally.


"A moment before adjournment, Dr. George B. Winter of St. Louis, retiring president of the association, surrendered the chair to Dr. Leroy M. S. Miner, incoming president, who will be succeeded next year by Dr. Camalier. Dr. Winter was guest of honor at the president's ball in the Hotel St. Francis . . .

"OFFICERS NAMED"

"Atlantic City, N. J., was chosen as the convention city for next year and the following officers were elected, their terms to commence immediately:

"Vice presidents: Dr. J. V. Gentilly, Cleveland, Ohio; Dr. J. A. Blue, Birmingham, Ala.

"The following were reelected: Secretary, Dr. Harry B. Pinney, Chicago; treasurer, Dr. R. H. Volland, Iowa City, Iowa; trustees, Dr. Harvey J. Burkhardt, Rochester, N. Y.; Dr. C. R. Lawrence, Enid, Okla.; Dr. Herbert C. Miller, Portland, Ore.; Dr. Wilfred H. Robinson, Oakland."—*From: San Francisco Chronicle, Friday, July 17, 1936.*



● Ask any dentist who is using Minimax Alloy No. 178 what his experience has been with Minimax. He will tell you that he has never had an occasion to feel sorry that he has used it.

Swing over to Minimax the next time you need alloy. Here is what you will get:

Alloy carefully designed to withstand the variables in amalgam technic over the wide range of manipulations present . . . so that whether the cavity is large or small, easily accessible or difficult to get at, it still retains the properties required by amalgam specifications!

It's the properties of amalgam that determine whether a filling will endure or fail. Look well to properties; swing over to Minimax.

The MINIMAX COMPANY

Medical & Dental Arts Bldg., Chicago, Ill.

MINIMAX

You Get More

Fillings suitable for alloy-mercury gauges.
Complies with Revised (1934) A.D.A. Specifications No. 1.

—You Pay Less

1 oz. bottle \$1.60
5 oz. bottle . . \$1.50 per oz.
10 oz. \$1.40 per oz.
20 oz. \$1.35 per oz.



IDEAL FOR THE GENERAL PRACTITIONER AND THE MAN WHO SPECIALIZES

Dental Oral Surgery really offers a postgraduate course in textbook form. It describes in complete detail the invaluable technique of the author, Doctor Wilton W. Cogswell. This is a book for the progressive man; it is ideal for both the general practitioner and the man who specializes. The price is \$10. Order your copy at once.

THE DENTAL DIGEST

1005 Liberty Ave.

Pittsburgh, Pa.

See page 257

D.D.8

WERNET DENTAL MFG. CO.
882 THIRD AVE., BROOKLYN, N. Y.

Send me free supply of Dr. Wernet's Powder as mentioned in ad.

Dr. _____

Address _____

City _____

See page 258

D.D.8

BRISTOL-MYERS CO.
19-S WEST 50TH STREET, NEW YORK CITY.

Please send sample of Sal Hepatica.

Dr. _____

Address _____

City _____

See page 277

D.D.8

EMERSON DRUG CO., BALTIMORE, MD.

Please send trial supply of Acetanilid as mentioned in ad.

Dr. _____

Address _____

City _____

See page 278

D.D.8

KELLY-BURROUGHS LABY., INC.
143 N. WABASH AVE., CHICAGO, ILL.

Send complete information concerning Kelly's Paste.

Dr. _____

Address _____

City _____

See page 279

D.D.8

WISCONSIN ALUMNI RESEARCH FOUND'T'N
MADISON, WISCONSIN

Please send 32 page booklet mentioned in ad.

Dr. _____

Address _____

City _____

See page 280

D.D.8

McKesson Appliance Co., Toledo, Ohio

Please send McKesson literature as mentioned in ad.

Dr. _____

Address _____

Dealer _____

See page 282

D.D.8


WILMOT CASTLE CO.
1109 UNIVERSITY AVE., ROCHESTER, N. Y.

Send me free booklet "Modern Sterilization" as mentioned in ad.

Dr. _____

Address _____

Dealer _____



WILSON'S
CO-RE-GA
(POWERED)
THE PERFECT
ADHESIVE for
DENTURES...

« « « Nothing excels the
proved effectiveness of
CO-RE-GA in helping
patients overcome the first
inhibitions of artificial
dentures » » » » »

PLEASE SEND FREE SAMPLES FOR PATIENTS

COREGA CHEMICAL CO.
201 ST. CLAIR AVE. N.W.
CLEVELAND OHIO, U.S.A.
This coupon is
for dentists
use only

MAIL COUPON FOR YOUR SUPPLY

CO-RE-GA is not advertised to the public



*CAL-C-MALT 'Roche'

Cal-C-Malt has remarkable tonic properties. It produces a sense of healthful euphoria, diminishes fatigability and improves appetite. There is no compensatory let-down following the invigorating effect of Cal-C-Malt, because Cal-C-Malt is not a stimulant.

Prescribe Cal-C-Malt to favor convalescence from febrile, infectious, or debilitating diseases; to overcome post-operative debility; as a nu-

trient for pregnant women, and also during the puerperium and the period of lactation; to patients of all ages, adults as well as children, who are undernourished.

In addition to its nutrient tonic properties Cal-C-Malt has important special indications because of its content of effective doses of calcium and vitamin C. Dose: Two teaspoonfuls in milk, t. i. d. *Package: 12-oz. tin.*

**Dibasic Calcium Phosphate and Vitamin C mixed with the finest quality wind-blown cocoa powder, and containing also vitamin B, iron, magnesium, and other minerals.*

Cal-C-Malt makes a delicious beverage — mix it with milk.

HOFFMANN-LA ROCHE • INC • NUTLEY • N. J.

See page 282

D.D.8

LEE S. SMITH & SON MFG. CO.

7325 PENN AVE., PITTSBURGH, PA.

Please send me a 3 oz. can of Foil Coat Thinner, for the regular retail price of \$1.90. I understand that you will send me a dollar can of Foil Coat Improved—free.

Dr. _____

Address _____

Dealer _____

See page 283

D.D.8

C. V. MOSBY COMPANY, 3525 PINE BLVD.
ST. LOUIS, MO.

Send me the following books: Nos. 1 (), 2 (), 3 (), 4 (), 5 (), 6 (), 7 (), charging my account. I agree to pay for these in accordance with your monthly payment plan.

Dr. _____

Address _____

City _____

See page 284

D.D.8

BUFFALO DENTAL MFG. CO.,
BUFFALO, N. Y.

Send me a \$1.00 jar of Buffalo Prophylaxis Paste through my dealer.

Dr. _____

Address _____

Dealer _____

See page 284

D.D.8

W. V-B. AMES CO., FREMONT, OHIO

Send me a copy of your booklet as mentioned in your ad, please.

Dr. _____

Address _____

City _____

See page 288

D.D.8

LEE S. SMITH & SON MFG. CO.,
7325 PENN AVE., PITTSBURGH, PA.

Send me 5 oz. Lee Smith Certified Alloy, with sufficient mercury for amalgamation, at your special price of \$8.60.

Dr. _____

Address _____

Dealer _____

See page 288

D.D.8

PELTON & CRANE COMPANY
632 HARPER AVE., DETROIT, MICH.

Send complete details concerning the new Pelton Localite. No obligation of course.

Dr. _____

Address _____

Dealer _____



Here is the Light which dentists have long demanded. Provides 300 foot-candles of cold, concentrated light *inside* the mouth, projected from any direction. Uses tiny 4-watt lamp with transformer built into supporting bracket, and generates no heat whatever. No glare in eyes of either operator or patient. Engineered by lighting specialists, and beautifully finished in chrome and black. Unit or Stand models, \$35.00. Wall bracket model, \$45.00. Complete details sent on request.

THE PELTON & CRANE COMPANY, DETROIT, MICHIGAN

THE NEW PELTON *Localite*

A Special Offer on

Lee Smith Certified Alloy

Here is an opportunity to make a real buy on alloy.

Five ounces Lee Smith Certified Alloy, with sufficient mercury for amalgamation, sold as a unit, \$8.60.

Lee Smith Certified Alloy represents the utmost in quality. As its name indicates, it is certified to exceed all A.D.A. specifications for amalgam alloys. And it will fit your alloy mercury proportioner.

Use the coupon on page 287 for convenience in ordering.

LEE S. SMITH & SON MFG. CO.
2725 PENN AVE., PITTSBURGH, PA.

SUGGESTIONS TO CONTRIBUTORS

PUBLICATION PREROGATIVE: Articles and illustrations are accepted for publication on condition that they are contributed solely to this magazine.

COPYRIGHT: All articles and illustrations appearing in *THE DENTAL DIGEST* are covered by copyright in the name of *THE DENTAL DIGEST, Incorporated*. Permission will be granted on request for the reproduction of text or illustrations in reputable publications (and other noncommercial purposes) of anything appearing in the pages of *THE DIGEST* if proper credit is given.

MANUSCRIPTS: Manuscripts should be typewritten, double-spaced, and the original, not the carbon copy submitted. Footnotes and bibliographies should have a definite connection with the article and should be specifically referred to in the article. To be of value to readers, bibliographic references should contain complete information in the order given: name of author, title of article, name of periodical, with volume, page, month—day of month if weekly—and year. In the case of books: name of author, title, edition, volume, place of publication, name of publishers, year, pages. Manuscripts should not be rolled.

ILLUSTRATIONS: Drawings and photographs should be plainly numbered on the backs according to the sequence in which reference is made to them in the article. The author's name should be written on the back of each illustration, and in case of possible doubt, the word "Top" to designate the top of the illustration. In the case of roentgenograms, the negatives are preferred for reproduction; in the case of photographs, glossy prints about 12 by 8 inches in size. Line drawings should be made in black on white paper that does not blot. Color work must be of a particularly high quality to be acceptable. All illustrations should be clear and distinct and large enough so that details will not be lost if reduction in size is necessary.

EDITING: Authors should not feel that they are being discriminated against or personally criticized when changes are made in the wording or spelling of their manuscripts or if parts are deleted. A minimum of editing is necessary in all cases—if for no other reason than to make grammatical corrections, and sometimes one article will require more revision than another. The reason for this is obvious. Every magazine has its peculiar style in matters of arbitrary spelling, in its general tone, in its form of presentation. The *Digest* favors a compact, terse, simple style, with outlining wherever possible, and many illustrations. Wordy, padded articles, with extraneous and irrelevant matter and florid writing will necessarily undergo considerable editing to make them conform to our style of succinct, *purposeful* writing. It is, however, at all times the aim of the editors to preserve the author's meaning and to help him make that meaning clear; the editing is not done for standardization as such.

ANONYMITY: Anonymous manuscripts and communications will not be read.



DECIDEDLY IMPORTANT

TO THOSE WHO CARE

Steele's
INTERCHANGEABLE
Tru pontics

Teeth and their importance to personal appearance have been stressed so often that today, more than ever before, patients demand and appreciate good bridgework.

The requirements of a perfect bridge tooth include strength, hygiene, esthetics and interchangeability. Steele's Tru pontics meet all these requirements.

Your patients will be quick to understand the advantages of a Tru pontic bridge. Use the best.

THE COLUMBUS DENTAL MFG. CO. Columbus, Ohio, U.S.A.



WHEN YOU SELECT SHADES FOR
GRAY BLONDES, NEW TRUBYTE 39,
41, 43, ARE MOST FREQUENTLY
REQUIRED.

NEW TRUBYTE SHADES
PROVIDE THE RIGHT COLORS IN
NATURAL BLENDINGS.

REMEMBER THERE ARE 24 USEFUL SHADES
IN NEW TRUBYTE.



THE DENTISTS' SUPPLY COMPANY OF NEW YORK